

Fig. 3.

Threshold E-field for producing burns in adult males (solid curves) and ten-year old children (dashed curves) in finger contact with various vehicles (contact area = 25 mm^2).




Fig. 4.

Threshold E-field for producing burns in adult males (solid curves) and ten-year old children (dashed curves) in finger contact with various vehicles (contact area = 144 mm²).

REFERENCES

1. O. P. Gandhi, I. Chatterjee, D. Wu and Y. G. Gu, "Likelihood of High Rates of Energy Deposition in the Human Legs at the ANSI Recommended 3-30 MHz RF Safety Levels," *Proceedings of IEEE*, Vol. 37, pp. 1145-1147, 1985.
2. D. A. Hill and J. A. Walsh, "Radio Frequency Current Through the Feet of a Grounded Human," *IEEE Transactions on Electromagnetic Compatibility*, Vol. 27, pp. 18-23, 1985.
3. O. P. Gandhi, J. Y. Chen and A. Riazzi, "Currents Induced in a Human Being for Plane Wave Exposure Conditions 0-50 MHz and for RF Sealers," *IEEE Transactions on Biomedical Engineering*, Vol. 33, pp. 757-767, 1986.
4. J. Y. Chen and O. P. Gandhi, "RF Currents Induced in an Anatomically-Based Model of a Human for Plane Wave Exposures 20-100 MHz," *Health Physics*, Vol. 57, pp. 89-98, 1989.
5. O. P. Gandhi, Y. G. Gu, J. Y. Chen and H. I. Bassen, "Specific Absorption Rates and Induced Current Distributions in an Anatomically Based Human Model for Plane Wave Exposures," *Health Physics*, Vol. 63, pp. 281-290, 1992.
6. O. P. Gandhi and J. F. DeFord, "Calculation of EM Power Deposition for Operator Exposure to RF Induction Heaters," *IEEE Transactions on Electromagnetic Compatibility*, Vol. 30, pp. 63-68, 1988.
7. J. Y. Chen and O. P. Gandhi, "RF Currents Induced in an Anatomically Based Model of Man for Leakage Fields of a Parallel Plate Dielectric Heater," *IEEE Transactions on Microwave Theory and Techniques*, Vol. 37, pp. 174-180, 1989.
8. J. Y. Chen, O. P. Gandhi and D. L. Conover, "SAR and Induced Current Distributions for Operator Exposure to RF Dielectric Sealers," *IEEE Transactions on Electromagnetic Compatibility*, Vol. 33, pp. 252-261, 1991.
9. C. F. Dalziel, and T. H. Mansfield, "Effect of Frequency on Perception Currents," *Trans. AIEE*, Vol. 69(II), pp. 1162-1168, 1950.
10. C. F. Dalziel, and W. R. Lee, "Lethal Electric Currents," *IEEE Spectrum*, Vol. 6, pp. 44-50, 1969.
11. R. J. Rogers, "Radio-Frequency Burn Hazards in the MF/HF Band," in *Aeromedical Review-Proc. Workshop on the Protection of Personnel Against RF Electromagnetic Radiation*, Review 3-81, J. C. Mitchell, Ed. (USAF School of Aerospace Medicine, Brooks Air Force Base, Texas 78235), pp. 76-89, Sept. 1981.
12. C. F. Dalziel, "The Threshold of Perception Currents," *Trans. AIEE*, Vol. 73, pp. 990-996, 1954.
13. C. F. Dalziel, "Electric Shock Hazard," *IEEE Spectrum*, Vol. 9, pp. 41-50, 1972.
14. O. P. Gandhi and I. Chatterjee, "Radio-Frequency Hazards in the VLF to MF Band," *Proceedings of the IEEE*, Vol. 70, pp. 1462-1464, 1982.
15. A. W. Guy and C. K. Chou, "Hazard Analysis: Very Low Frequency Through Medium Frequency Range," Final report USAF SAM Contract No. F 33615-78-D-0617, Task 0065, 1982.

16. A. W. Guy and C. K. Chou, "Very Low Frequency Hazard Study," Final report prepared for USAF School of Aerospace Medicine, Brooks Air Force Base, TX, Contract No. F 33615-83-C-0625, 1985.
17. I. Chatterjee, D. Wu and O. P. Gandhi, "Human Body Impedance and Threshold Currents for Perception and Pain for Contact Hazard Analysis in the VLF-MF Band," *IEEE Transactions on Biomedical Engineering*, Vol. 33, pp. 486-494, 1986.
18. D. W. Deno and L. E. Zaffanella, "Electrostatic Effects of Overhead Transmission Lines and Stations," Chapter 8 in *Transmission Line Reference Book: 345 kV and Above*, 2nd Edition, published by Electric Power Research Institute, Palo Alto, CA, 1982.
19. J. P. Reilly, "Electrical Stimulation and Electropathology," Cambridge University Press, 1992.
20. ANSI/IEEE C95.1-1992, "Standard for Safety Levels with Respect to Human Exposure to Radio Frequency Electromagnetic Fields, 3 kHz to 300 GHz," available from the Institute of Electrical and Electronics Engineers, Inc., 345 East 47th Street, New York, NY 10017.
21. O. P. Gandhi, I Chatterjee, D. Wu, J. A. D'Andrea and K. Sakamoto, "Very Low Frequency (VLF) Hazard Study," Final report USAF SAM Contract No. F33615-83-R-0613, 1985.
22. O. P. Gandhi, "ANSI Radiofrequency Safety Guide: Its Rationale, Some Problems and Suggested Improvements," Chapter 3 in *Biological Effects and Medical Applications of Electromagnetic Energy*, O. P. Gandhi, Ed., Prentice Hall Inc., 1990.
23. O. P. Gandhi, "Basis for RFR-Safety Standards in the 10 kHz-50 MHz Region," Final Report submitted to USAF SAM Contract No. F 33615-85-R-4522, Brooks Air Force Base, TX, 1987.
24. L. P. Ferris, B. G. King, P. W. Spence, and H. B. Williams, "Effect of Electric Shock on the Heart," *Trans. AIEE*, Vol. 55, pp. 498-515, 1936.
25. C. F. Dalziel, "Dangerous Electric Currents," *Trans. AIEE*, Vol. 65, pp. 579-585, 1946.
26. C. F. Dalziel, "Effects of Electric Shock on Man," *IRE Trans. Medical Electronics*, PGME-5, pp. 44-62, 1956.
27. W. R. Lee, "Death From Electric Shock," *Proc. IEE*, Vol. 113, pp. 144-148, 1966.
28. M. S. Hammam and R. S. Baishiki, "A Range of Body Impedance Values for Low Volume, Low Source Impedance Systems of 60 Hz," *IEEE Trans. Power Apparatus and Systems*, PAS-102, pp. 1097-1103, 1983.
29. L. A. Geddes, L. E. Baker, P. Cabler, and D. Brittain, "Response to Passage of Sinusoidal Current Through the Body," In, N. L. Wulfsohn and A. Sances (eds.), *The Nervous System and Electric Current*, Vol. 2, New York, Plenum Press, pp. 121-129, 1971.
30. C. M. Becker, I. V. Malhotra, and J. Hedley-Whyte, "The Distribution of Radio Frequency Current and Burns," *Anesthesiology*, Vol. 38, pp. 106-122, 1973.
31. A. K. Dobbie, "The Electrical Aspects of Surgical Diathermy," *Biomedical Engineering*, Vol. 4, pp. 206-216, 1969.

32. L. R. Delaplace and J. P. Reilly, "Electric and Magnetic Field Coupling from High Voltage AC Power Transmission Lines -- Classification of Short-Term Effects on People," *IEEE Transactions on Power Apparatus and Systems*, Vol. PAS-97, pp. 2243-2252, Nov./Dec. 1978.

E203

NOTICE: This material may be protected
by copyright law (Title 17 U.S. Code)

THE ENVIRONMENTAL HEALTH CRITERIA SERIES

- Acrolein (No. 127, 1991)
- Acrylamide (No. 49, 1985)
- Acrylonitrile (No. 28, 1983)
- Aldicarb (No. 121, 1991)
- Aldrin and dieldrin (No. 91, 1989)
- Allethrin (No. 87, 1989)
- Ammonia (No. 54, 1986)
- Arsenic (No. 18, 1981)
- Asbestos and other natural mineral fibres (No. 53, 1986)
- Barium (No. 107, 1990)
- Beryllium (No. 106, 1990)
- Biotoxins, aquatic (marine and freshwater) (No. 37, 1984)
- Butanols - four isomers (No. 65, 1987)
- Cadmium (No. 134, 1992)
- Cadmium - environmental aspects (No. 135, 1992)
- Camphechlor (No. 45, 1984)
- Carbamate pesticides: a general introduction (No. 64, 1986)
- Carbon disulfide (No. 10, 1979)
- Carbon monoxide (No. 13, 1979)
- Carcinogens, summary report on the evaluation of short-term *in vitro* tests (No. 47, 1985)
- Carcinogens, summary report on the evaluation of short-term *in vivo* tests (No. 109, 1990)
- Chlordane (No. 34, 1984)
- Chlordecone (No. 43, 1984)
- Chlorine and hydrogen chloride (No. 21, 1982)
- Chlorobenzenes other than hexachlorobenzene (No. 128, 1991)
- Chlorofluorocarbons, fully halogenated (No. 113, 1990)
- Chlorofluorocarbons, partially halogenated (methane derivatives) (No. 126, 1991)
- Chlorophenols (No. 93, 1989)
- Chromium (No. 61, 1988)
- Cyhalothrin (No. 99, 1990)
- Cypermethrin (No. 82, 1989)
- 1,2-Dichloroethane (No. 62, 1987)
- 2,4-Dichlorophenoxyacetic acid (2,4-D) (No. 29, 1984)
- 2,4-Dichlorophenoxyacetic acid - environmental aspects (No. 84, 1989)
- DDT and its derivatives (No. 9, 1979)
- DDT and its derivatives - environmental aspects (No. 83, 1989)
- Deltamethrin (No. 97, 1990)
- Diaminotoluenes (No. 74, 1987)
- Dichlorvos (No. 79, 1988)
- Diethylhexyl phthalate (No. 131, 1992)
- Dimethoate (No. 90, 1989)
- Dimethylformamide (No. 114, 1991)
- Dimethyl sulfate (No. 48, 1985)
- Diseases of suspected chemical etiology and their prevention, principles of studies on (No. 72, 1987)
- Dithiocarbamate pesticides, ethylenethiourea, and propylenethiourea: a general introduction (No. 78, 1988)
- Endosulfan (No. 40, 1984)
- Endrin (No. 130, 1992)
- Environmental epidemiology, guidelines on studies in (No. 27, 1983)
- Epichlorohydrin (No. 33, 1984)
- Ethylene oxide (No. 55, 1985)
- Extremely low frequency (ELF) fields (No. 35, 1984)
- Fenitrothion (No. 133, 1992)
- Fenvalerate (No. 95, 1990)
- Fluorine and fluorides (No. 36, 1984)
- Food additives and contaminants in food, principles for the safety assessment of (No. 70, 1987)
- Formaldehyde (No. 89, 1989)
- Genetic effects in human populations, guidelines for the study of (No. 46, 1985)
- Heptachlor (No. 38, 1984)
- Alpha- and beta-hexachlorocyclohexanes (No. 123, 1991)
- Hexachlorocyclopentadiene (No. 120, 1991)
- n-Hexane (No. 122, 1991)
- Hydrazine (No. 68, 1987)
- Hydrogen sulfide (No. 19, 1981)
- Infancy and early childhood, principles for evaluating health risks from chemicals during (No. 59, 1986)
- Isobenzan (No. 129, 1991)
- Kelevan (No. 66, 1986)
- Lasers and optical radiation (No. 23, 1982)
- Lead (No. 3, 1977)*

This report contains the collective views of an international group of experts and does not necessarily represent the decisions or the stated policy of the United Nations Environment Programme, the International Radiation Protection Association, or the World Health Organization.

Environmental Health Criteria 137

ELECTROMAGNETIC FIELDS (300 Hz to 300 GHz)

Published under the joint sponsorship of
the United Nations Environment Programme,
the International Radiation Protection Association,
and the World Health Organization



World Health Organization
Geneva, 1993

* Out of print

continued inside back cover

WHO Library Cataloguing in Publication Data

Electromagnetic fields (300 Hz to 300 GHz)

(Environmental health criteria ; 137)

1. Electromagnetic fields - adverse effects 2. Environmental exposure I. Series

ISBN 92 4 157137 3 (NLM Classification: QT 34)
ISSN 0250-863X

The World Health Organization welcomes requests for permission to reproduce or translate its publications, in part or in full. Applications and enquiries should be addressed to the Office of Publications, World Health Organization, Geneva, Switzerland, which will be glad to provide the latest information on any changes made to the text, plans for new editions, and reprints and translations already available.

©World Health Organization 1993

Publications of the World Health Organization enjoy copyright protection in accordance with the provisions of Protocol 2 of the Universal Copyright Convention. All rights reserved.

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the Secretariat of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

Printed in Finland
92/9531 - Vammala - 5500

CONTENTS

PREFACE	11
1. SUMMARY AND RECOMMENDATIONS FOR FURTHER STUDIES	15
1.1 Summary	15
1.1.1 Physical characteristics in relation to biological effects	15
1.1.2 Sources and exposure	16
1.1.2.1 Community	16
1.1.2.2 Home	16
1.1.2.3 Workplace	16
1.1.3 Biological effects	17
1.1.4 Laboratory studies	18
1.1.5 Human studies	20
1.1.6 Health hazard assessment	21
1.1.6.1 Thermal effects	21
1.1.6.2 Pulsed fields	22
1.1.6.3 Amplitude-modulated RF fields	22
1.1.6.4 RF field effects on tumour induction and promotion	22
1.1.6.5 RF-induced current densities	22
1.1.6.6 RF contact shocks and burns	22
1.1.7 Exposure standards	23
1.1.7.1 Basic exposure limits	23
1.1.7.2 Occupational exposure limits	24
1.1.7.3 Exposure limits for the general population	25
1.1.7.4 Implementation of standards	25
1.1.8 Protective measures	25
1.2 Recommendations for further studies	26
1.2.1 Introduction	26
1.2.2 Pulsed fields	26
1.2.3 Cancer, reproduction, and nervous system studies	27
1.2.4 Weak-field interactions	27
1.2.5 Epidemiology	28
2. PHYSICAL CHARACTERISTICS	29
2.1 Introduction	29

2.2 Electric field	29
2.3 Magnetic field	31
2.4 Waves and radiation	32
3. NATURAL BACKGROUND AND HUMAN-MADE SOURCES	36
3.1 General	36
3.2 Natural background	37
3.2.1 Atmospheric fields	37
3.2.2 Terrestrial emissions	38
3.2.3 Extraterrestrial fields	38
3.3 Human-made sources	39
3.3.1 General	39
3.3.2 Environment, home, and public premises	43
3.3.3 Workplace	45
3.3.4 Medical practice	48
4. EXPOSURE EVALUATION - CALCULATION AND MEASUREMENT	51
4.1 Introduction	51
4.2 Theoretical estimation	51
4.3 Measurements	56
4.3.1 Preliminary considerations	56
4.3.2 Near-field versus far-field	57
4.3.3 Instrumentation	57
4.3.4 Measurement procedures	58
5. DOSIMETRY	61
5.1 General	61
5.2 Low frequency range	62
5.2.1 Magnetic fields	62
5.2.2 Electric fields	65
5.3 High-frequency range	72
5.4 Derivation of exposure limits from basic quantities	78
6. INTERACTION MECHANISMS	80
6.1 General	80
6.2 Electrical properties of cells and tissues	81
6.2.1 Permittivity	81
6.2.2 Non-linear effects	84
6.2.3 Induced fields at the cellular level	85

6.2.4 Body impedance	86
6.3 Direct interactions - strong fields	88
6.3.1 Interactions with excitable tissues	88
6.3.2 Thermal interactions	89
6.4 Direct interactions - weak fields	91
6.4.1 General	91
6.4.2 Microelectrophoretic motion	93
6.4.3 Ion-resonance conditions	93
6.4.4 Calcium ion exchange	94
6.5 Indirect interactions	96
7. CELLULAR AND ANIMAL STUDIES	104
7.1 Introduction	104
7.2 Macromolecules and cell systems	104
7.2.1 Effects on cell membranes	105
7.2.2 Effects on haematopoietic tissue	107
7.2.3 Isolated cerebral tissue, peripheral nerve tissue, and heart preparations	110
7.2.4 Mutagenic effects	111
7.2.5 Cancer-related studies	112
7.2.6 Summary and conclusions: <i>in vitro</i> studies	114
7.3 Animal studies	114
7.3.1 Nervous system	115
7.3.2 Ocular effects	118
7.3.3 Auditory perception	123
7.3.4 Behaviour	124
7.3.4.1 Thermoregulation	124
7.3.4.2 Activity (spontaneous movement)	126
7.3.4.3 Learned behaviours	127
7.3.5 Endocrine system	130
7.3.6 Haematopoietic and immune systems	133
7.3.7 Cardiovascular system	139
7.3.8 Reproduction and development	141
7.3.8.1 kHz studies	141
7.3.8.2 MHz and GHz studies	142
7.3.9 Genetics and mutagenesis	146
7.3.10 Cancer-related studies	148
7.3.11 Summary and conclusions	152
8. HUMAN RESPONSES	155
8.1 Laboratory studies	155
8.1.1 Cutaneous perception	155

WHO/IRPA TASK GROUP ON ELECTROMAGNETIC FIELDS (300 Hz TO 300 GHz)

Members

Prof J. Bernhardt ^a	Federal Office of Radiological Protection, Institute for Radiation Hygiene, Munich- Neuherberg, Germany
Dr C. F. Blackman	US Environmental Protection Agency, Health Effects Research Laboratory, North Carolina, USA
Dr L.A. Court ^a	Département de Protection Sanitaire, Centre d'Etudes Nucléaires, Fontenay-aux-Roses, France
Mme A. Duchêne ^a	Scientific Secretary, International Non-ionizing Radiation Committee, Fontenay-aux-Roses, France
Prof M. Grandolfo ^a	Istituto Superiore di Sanità, Rome, Italy
Dr M.H. Repacholi ^a	Royal Adelaide Hospital, Adelaide, Australia (<i>Chairman</i>)
Dr R.D. Saunders	National Radiological Protection Board, Didcot, United Kingdom (<i>Co-Rapporteur</i>)
Prof M.G. Shandala ^a	AN Marzeev Research Institute of General and Communal Hygiene, Kiev, USSR
Dr J.A. Stolwijk ^a	Department of Epidemiology and Public Health, Yale University, New Haven, USA
Dr M.A. Stuchly ^a	Bureau of Radiation and Medical Devices, Ottawa, Canada

8.1.2 Other perception thresholds	158
8.1.3 Auditory effects	158
8.1.4 Induced-current effects	158
8.1.5 Thermoregulation	159
8.1.6 Contact currents	164
8.2 Epidemiological and clinical comparative studies	164
8.2.1 Mortality and morbidity studies	164
8.2.2 Ocular effects	168
8.2.3 Effects on reproduction	168
8.2.4 VDU studies	171
8.2.5 Conclusions	173
8.3 Clinical case studies and accidental overexposures	173
9. HEALTH HAZARD ASSESSMENT	175
9.1 Introduction	175
9.2 Thermal effects	176
9.3 RF contact shocks and burns	177
9.4 Induced current densities	178
9.5 Pulsed RF fields	179
9.6 RF fields amplitude modulated at ELF frequencies	179
9.7 RF effects on tumour induction and progression	180
10. EXPOSURE STANDARDS	181
10.1 General considerations	181
10.2 Present trends	182
10.3 Recommendations by the IRPA	183
10.4 Concluding remarks	186
11. PROTECTIVE MEASURES	187
11.1 Engineering measures	187
11.2 Administrative controls	190
11.3 Personal protection	191
11.4 Medical surveillance	191
11.5 Interference with medical devices and safety equipment	193
GLOSSARY	194
REFERENCES	206
RESUME ET RECOMMANDATIONS EN VUE D'ETUDES FUTURES	259
RESUMEN Y RECOMENDACIONES PARA ESTUDIOS ULTERIORES	276

Dr M. Swicord Centre for Devices and Radiological
Health, Food and Drug
Administration, Rockville, USA
(*Co-Rapporteur*)

Dr L.D. Szabo^a National Research Institute for
Radiobiology and Radiation Hygiene,
Budapest, Hungary

Dr S. Szmigielski Centre for Radiobiology and
Radiation Safety, Warsaw, Poland
(*Vice-Chairman*)

Observer

Dr A. McKinlay^a National Radiological Protection
Board, Didcot, United Kingdom

^aFrom the International Non-Ionizing Radiation Committee of IRPA.

NOTE TO READERS OF THE CRITERIA MONOGRAPHS

Every effort has been made to present information in the criteria monographs as accurately as possible without unduly delaying their publication. In the interest of all users of the environmental health criteria monographs, readers are kindly requested to communicate any errors that may have occurred to the Director of the International Programme on Chemical Safety, World Health Organization, Geneva, Switzerland, in order that they may be included in corrigenda, which will appear in subsequent volumes.

frequencies (assessed as the decreased survival of implanted embryos and fetuses). Much experimental evidence suggests that acute or long-term RF exposures do not result in an increase in chromosome aberration frequency, when temperatures are maintained within physiological limits. One study reported an increased frequency of cytogenetic effects in mice exposed long-term at SARs between 0.05 and 20 W/kg. However, this study was not successfully corroborated using a different strain of mouse.

In general, the data in Table 28 suggest that the only exposures that are potentially mutagenic are those at high RF power densities, which result in substantial increase in temperature.

7.3.10 Cancer-related studies

A summary of cancer-related animal studies is given in Table 29. The number and types of studies are limited.

Exposure to RF levels sufficiently high to induce hyperthermia has generally resulted in tumour regression following transplantation of tumour cells (Preskorn et al., 1978; Roszkowski et al., 1980). In contrast, an increase in tumour progression has been observed in mice exposed long-term at lower, possibly thermogenic, SARs (Szmigielski et al., 1982). This effect was related to a non-specific stress. The authors suggested a transient shift in immune surveillance resulting in a lowering of resistance to neoplastic growth, as a likely explanation. Exposure at about 1 W/kg did not have any effect on melanoma growth in mice (Santini et al., 1988).

The effects of exposure on spontaneous or chemically-induced tumours have also been examined. In contrast to transplantation studies, these can test for an effect on the process of carcinogenesis. Two early studies (Prausnitz & Suskind, 1962; Skidmore & Baum, 1974), relevant to cancer induction, but in which the methodology was flawed in relation to an analysis of this end-point, are described for completeness. An increased incidence of monocytic leukaemia (defined as a non-circulating neoplasm of white-blood cells) and lymphatic or myeloid leukaemia (defined as a circulating "leukosis") was reported in Swiss mice exposed to thermally significant levels (half the acute LD₅₀) of 9.27 GHz pulsed RF, for 5 days per week

Table 29. Cancer-related studies

Exposure conditions	Effect on exposed group	Reference
Transplanted tumour cells		
2.45 GHz (CW), 35 W/kg, for 20 min/day during days 11-14 of gestation; offspring injected with sarcoma cells at 16 days of age exposed for 36 days	Retarded tumour growth and tumour incidence in sarcoma-injected offspring of exposed pregnant mice; rectal temperature of dams rose over 2 °C; exposed mice had increased longevity	Preskorn et al. (1978)
2.45 GHz (CW), 25 W/kg, 2 h/day for 7 days; Injection of sarcoma cells in mice 14 days after, or just after, RF exposure	Temporary tumour regression followed by renewed tumour growth 12 days later, when exposure 14 days after tumour injection; accelerated tumour growth, if exposed before implantation of tumour; lung metastases increased	Roszkowski et al. (1980)
2.45 GHz (CW), 2-3 W/kg or 6-8 W/kg, 2 h/day, for 6 days/week; mice exposed from 6 weeks of age to 12 months of stress	RF caused increase in sarcoma colonies in lungs in mice injected intravenously with these cells; chronic via confinement caused similar increase in lung tumours as 2-3 W/kg, but 6-8 W/kg produced higher increase in tumours	Szmigielski et al. (1982)
2.45 GHz (CW and pulsed) 10 W/m ² , 1.2 W/kg prior to, and during, B16 melanoma tumour transplantation and growth; exposed for 2.5 h/day, 6 times/week for 15 days, prior to injection of melanoma cells, then exposed to same schedule until death	No difference in mean tumour surface area/animal, or in mean survival time between exposed or control mice	Santini et al. (1988)
Spontaneous or chemically-induced tumours		
2.45 GHz (CW), 2-3 W/kg or 6-8 W/kg, 2 h/day, for 6 days/week, mice exposed from 6 weeks of age to 12 months of stress	SAR-dependent acceleration of mammary tumours in mice genetically predisposed to these tumours, and acceleration of skin tumours in mice painted with the carcinogen 3,4-benzopyrene (BP)	Szmigielski et al. (1982)

Table 29 (continued)

Exposure conditions	Effect on exposed group	Reference
2.45 GHz (CW), 100 W/m ² 4-5 W/kg, for 2 h/day, 5-6 days/week for a few months	Increased development of chemically-induced hepatomas and sarcomas in mice; survival of exposed mice decreased; increased frequency of skin tumours in mice given subcarcinogenic dose of BP	Szmigielski et al. (1988)
2.45 GHz (10 μ s pulses at 800 Hz) square wave- modulated at 8 Hz, 0.4 W/kg, continuous exposure at 2-27 months of age (lifetime study of rats)	Total incidence of neoplasia not significantly different from that in controls; however, increased number of primary malignancies (18) occurred early in exposed group compared with controls (5)	Guy et al. (1985)

for 59 days (Prausnitz & Susskind, 1962). However, the study suffered several deficiencies: leukosis and leukaemia were inadequately defined, infection may well have confounded the results, a large proportion of mice died without a cause of death being identified, and statistical analysis was absent (Roberts 1983; Kirk 1984).

Skidmore & Baum (1974) reported that exposure for 5 days per week for 33 weeks to very short pulses (5 ns rise time; 550 ms decay time) of high field strength (447 kV/m) pulsed at 5 Hz, resulted in a reduced incidence of leukaemia in AKR/J mice (which spontaneously develop a high incidence of lymphatic leukaemia between 26 and 52 weeks of age) compared with controls at the end of the exposure. However, the absence of a complete analysis of leukaemia incidence (and other causes of death) precludes any conclusion being drawn from this study. The authors also reported a zero incidence of mammary tumours in 1-year-old female Sprague-Dawley rats that had been exposed for 38 weeks; evaluation was probably premature for this end-point, the tumours occur spontaneously mainly in older rats. A later study (Baum et al., 1976) reported no effects on mammary tumour incidence and other lesions in rats exposed for 94 weeks.

Two studies merit particular attention. The long-term exposure of mice at SARs of between 2 and 8 W/kg resulted in an increase in

the number of sarcoma cell colonies in the lungs (following the injection of sarcoma cells), as shown in Fig. 22, and in an SAR-dependent increase in the rate of development of spontaneous mammary tumours and chemically-induced skin tumours. Repeated microwave exposure, followed by a "sub-carcinogenic" dose of carcinogen, resulted in an increased number of skin tumours. A study of 100 rats exposed for most of their lifetime at about 0.4 W/kg did not show any increased incidence of non-neoplastic lesions compared with control animals; longevity was very similar in both groups. However, the overall incidence of primary malignancy in the exposed group (18) was significantly greater than the control




Fig. 22. Number of lung tumours (following intravenous injection of 2×10^5 viable sarcoma cells) in mice exposed to 2.45 GHz microwaves or non-specific stress (overcrowding; Oc). From Szmigielski et al. (1988).

value (5), but was reported to be similar to the spontaneous incidence given in the literature for the particular strain of rat. Under these circumstances, it is difficult to draw any firm conclusions.

Tumour weights were not significantly different in rats implanted with mammary adenocarcinoma tissue and either exposed 25 days later to 2 kHz magnetic fields of up to 2 mT for 1 h a day for 9 days or not exposed (Baumann et al., 1989). Handling and restraint stress in animals were identified as possible confounders for the detection of subtle magnetic field effects.

7.3.11 Summary and conclusions

Most of the biological effects of acute exposure to RF fields are consistent with responses to induced heating, resulting either in rises in tissue or body temperature of about 1 °C or more, or in responses for minimizing the total heat load. Most responses in different animal species, exposed under various environmental conditions, have been reported at SARs above about 1-2 W/kg.

These animal (particularly primate) data indicate the types of response that are likely to occur in humans subject to a sufficient heat load. However, direct quantitative extrapolation to humans is difficult, given species differences in responses, in general, and in thermoregulatory ability particularly.

The most sensitive animal responses to heat loads are thermoregulatory adjustments, such as reduced metabolic heat production and vasodilation, with thresholds ranging between about 0.05 and 5 W/kg, depending on environmental conditions. However, these reactions form part of the natural repertoire of thermoregulatory responses that serve to maintain normal body temperatures.

Transient effects seen in exposed animals that are consistent with responses to increases in body temperature of 1 °C or more (and/or SARs in excess of about 2 W/kg in primates and rats) include the reduced performance of learned tasks and increased plasma corticosteroid levels. Other heat-related effects include temporary haematopoietic and immune responses, possibly in conjunction with elevated corticosteroid levels. The most consistent effects observed are reduced levels of circulating lymphocytes and increased levels of

neutrophils, decreased natural killer cell function, and increased macrophage activation; an increase in the primary antibody response of B-lymphocytes has also been reported. Cardiovascular changes consonant with increased heat load, such as increased heart rate and cardiac output, have been observed, together with a reduction in the effects of drugs, such as barbiturates, the action of which can be altered by changes in circulation and clearance rates.

Most animal data indicate that implantation and the development of the embryo and fetus are unlikely to be affected by exposures that increase maternal body temperature by less than 1 °C. Above these temperatures, adverse effects, such as losses in implantation, growth retardation, and post-natal changes in behaviour, may occur, with more severe effects occurring at higher maternal temperatures.

Most animal data suggest that low RF exposure that does not raise body temperatures above the normal physiological range is not mutagenic; thus, such exposure will not result in somatic mutation or hereditary effects.

There is much less information describing the effects of long-term, low-level exposure. So far, it is not apparent that any long-term adverse effects can result from exposures below thermally significant levels. The animal data indicate that male fertility is unlikely to be affected by long-term exposure at levels insufficient to raise body and testis temperatures. Cataracts have not been induced in rabbits exposed at 100 W/m² for 6 months, or in primates exposed at 1.5 kW/m² for 3 months.

A study of 100 rats, exposed for most of their lifetime at about 0.4 W/kg, did not show an increased incidence of non-neoplastic lesions or total neoplasias compared with control animals; longevity was very similar in both groups. There were differences in the overall incidence of primary malignancies, but these could not necessarily be attributed to the RF exposure. The possibility that exposure to RF might influence the process of carcinogenesis is of particular concern. So far, there is no definite evidence that RF exposure does have an effect, but there is clearly a need for further studies to be carried out. Overwhelmingly, the experimental data indicate that RF fields are not mutagenic, and so they are unlikely to act as initiators of carcinogenesis. In a few studies, evidence has been sought of an enhancement of the effect of a known carcinogen.

The long-term exposure of mice at 2-8 W/kg resulted in an increase in the progression of spontaneous mammary tumours and of skin tumours in mice the skin of which was tested with a chemical carcinogen. Repeated RF exposure followed by a "sub-carcinogenic" dose of carcinogen resulted in an increased number of skin tumours; however, this study has been reported only briefly, and the authors noted the need for experimental confirmation.

In *in vitro* studies, enhanced cell transformation rates were reported after RF exposure at 4.4 W/kg (alone or combined with X-radiation) followed by treatment with a chemical promotor. The latter data have not always been consistent between studies. It is clear that studies relevant to carcinogenesis need replicating and extending further, to reduce uncertainties in this area.

A substantial body of data exists describing *in vitro* biological responses to amplitude-modulated RF radiation at SARs too low to involve any response to heating. Some studies have reported effects after exposure at SARs of less than 0.01 W/kg, occurring within modulation frequency "windows" (usually between 1 and 100 Hz) and sometimes within power density "windows".

Changes have been reported in the electroencephalograms of cats and rabbits, in calcium ion mobility in the brain tissue *in vitro* and *in vivo*, in lymphocyte cytotoxicity *in vitro*, and in the *in vitro* activity of an enzyme involved in cell growth and division. Some of these responses have been difficult to confirm, and their physiological or pathological consequences are not clear. However, any toxicological investigation should be based on tests carried out at appropriate levels of exposure. It is important that these studies be confirmed and extended to *in vivo* studies and that the health implications, if any, for exposed people are determined. Of particular importance, would be studies that link extremely low frequency, amplitude-modulated RF interactions at the cell surface with changes in DNA synthesis or transcription. It is worth noting that this interaction implies a "demodulation" of the RF signal at the cell membrane.

8. HUMAN RESPONSES

Epidemiology can be defined as the study of the occurrence of illness; its main goals are to evaluate hypotheses about the causation of illness and to relate disease occurrence to the characteristics of people and their environment. Epidemiological studies of human populations exposed to RF fields are few in number and are generally limited in scope. The principal groups studied have been people occupationally exposed in the military or in industry. Information about worker health status has generally come from medical records, questionnaires, and physical and laboratory examinations. Exposure data have come from personnel records, questionnaires, environmental measurements, and equipment-emission measurements. Determination of actual exposure to RF fields and to other risk factors for the same outcome is difficult in retrospective human studies.

Some studies of controlled exposures of volunteers have provided valuable information on responses to RF exposure. These studies include warming and pain thresholds for RF heating of the skin, RF hearing, and RF shocks and burns. Clinical studies of accidental overexposures provide information on acute-exposure responses.

8.1 Laboratory studies

8.1.1 Cutaneous perception

Exposure of the human body to RF fields can cause heating that is detectable by the temperature-sensitive receptors in the skin. Several investigators have determined experimentally the threshold intensities that cause sensations of perceptible warmth, pain, and delay in response to the stimulus in human subjects, as shown in Table 30.

Adair (1983a) noted that RF exposures to frequencies of 30 GHz and above would probably be similar to infrared in their perception threshold values. However, over much of the RF spectrum, current standards are set at levels that are below those that most would consider detectable by sensation. Thus, cutaneous perception may be an indicator of exposure only at RF frequencies of the order of several gigahertz or more, which have wavelengths that are small in comparison with the length of the exposed body, i.e., wavelengths

- GRANDOLFO, M., MICHAELSON, S.M., & RINDI, A., ed. (1983) *Biological effects and dosimetry of nonionizing radiation: radiofrequency and microwave energies*. New York, London, Plenum Press, p. 669.
- GRANDOLFO, M., VECCHIA, P., & GANDHI O.P. (1990) Magnetic resonance imaging calculation of radiofrequency power deposition in the human torso model. *Bioelectromagnetics*, 11: 117-128.
- GRUNDLER, W. & KEILMANN, F. (1983) Sharp resonances in yeast growth prove nonthermal sensitivity to microwaves. *Phys. Rev. Lett.* 51(13): 1214-1216.
- GRUNDLER, W. & KEILMANN, F. (1989) Resonant microwave effect on locally fixed yeast microcolonies. *Z. Naturforsch. (C)*, 44(9-10): 863-866.
- GUY, A.W. (1985) Hazards of VLF electromagnetic fields. In: *The impact of proposed radiofrequency radiation standards on military operations*. Neuilly-sur-Seine, France, Advisory Group for Aerospace Research and Development (AGARD), pp. 9.1-9.20 (Lecture Series No. 138).
- GUY, A.W. (1987) Dosimetry associated with exposure to nonionizing radiation: very low frequency to microwaves. *Health Phys.*, 53: 569-584.
- GUY, A.W. & CHOU, C.K. (1982) Hazard analysis: Very low frequency through medium frequency range, Texas, Brooks Air Force Base, USAF School of Aerospace Medicine, Aerospace Medical Division (Report USAFSAM 33615-78-D-0617).
- GUY, A.W., CHOU, C.K., LIN, J.C., & CHRISTENSEN, D. (1975a) Microwave-induced effects in mammalian auditory systems and physical materials. *Ann. N.Y. Acad. Sci.*, 247: 194-218.
- GUY, A.W., LIN, J.C., KRAMAR, P.O., & EMERY, A.F. (1975b) Effect of 2450 MHz radiation on the rabbit eye. *IEEE Trans. microwave Theory Tech.*, MTT-23: 492-498.
- GUY, A.W., KRAMAR, P.O., HARRIS, C.A., & CHOU, C.K. (1980) Long-term 2450 MHz CW microwave irradiation of rabbits: Methodology and evaluation of ocular and physiologic effects. *J. microwave Power*, 15: 37-44.
- GUY, A.W., DAVIDOW, S., YUANG, G.Y., & CHOU, C.K. (1982) Determination of electric current distributions in animals and humans

- exposed to a uniform 60-Hz high-intensity electric field. *Bioelectromagnetics*, 3: 47-71.
- GUY, A.W., CHOU, C.K., & NEUHAUS, B. (1984) Average SAR and SAR distribution in man exposed to 450 MHz radiofrequency radiation. *IEEE Trans. microwave Theory Tech.*, MTT-32: 752-762.
- GUY, A.W., CHOU, C.K., KUNZ, L.L., CROWLEY, J., & KRUPP, J. (1985) Effects of long-term low-level radiofrequency radiation exposure on rats. Volume 9. Summary. Texas, Brooks Air Force Base, USAF School of Aerospace Medicine (USFSAM-TR-85-11).
- GUY, A.W., CHOU, C.K., McDOUGALL, J.A., & SORENSEN, C. (1987) Measurement of shielding effectiveness of microwave-protective suits. *IEEE Trans. microwave Theory Tech.*, 35: 984-993.
- HAGMANN, M.J., LEVIN, R.L., & TURNER, P.F. (1985) A comparison of the annular phased array to helical coil applicators for limb and torso hyperthermia. *IEEE Trans.*, BME-32: 916-927.
- HALL, A. & BURSTOW, D.J. (1980) Risk of ignition of flammable gases and vapours by radio transmission. *Electrotechnology*, Jan: 12-15.
- HALLE, B. (1988) On the cyclotron resonance mechanism for magnetic field effects on transmembrane ion conductivity. *Bioelectromagnetics*, 9(4): 381-385.
- HAMRICK, P.E. & FOX, S.S. (1977) Rat lymphocytes in cell culture exposed to 2450 MHz (CW) microwave radiation. *J. microwave Power*, 12: 125-132.
- HAMRICK, P.E. & ZINKL, J.G. (1975) Exposure of rabbit erythrocytes to microwave irradiation. *Radiat. Res.*, 62: 164.
- HAMBURGER, S., LOGUE, J.N., & STERNTHAL, P.M. (1983) Occupational exposure to non-ionizing radiation and an association with heart disease: an exploratory study. *J. chronic Dis.*, 36: 791-802.
- HANKIN, N.N. (1974) An evaluation of selected satellite communications systems as sources of environmental microwave radiation. Silver Springs, Maryland, US Environmental Protection Agency (Report 520/2-74-008).
- HARVEY, S.M. (1984) Electric-field exposure of persons using video display units. *Bioelectromagnetics*, 5: 1-12.

References

- O'CONNOR, M.E. (1980) Mammalian teratogenesis and radio-frequency fields. *Proc. IEEE*, **68**: 56-60.
- ODLAND, L.T. (1973) Radiofrequency energy: A hazard to workers? *Ind. Med. Surg.*, **42**: 23-26.
- OLCERST, R.B., BELMAN, S., EISENBUD, M., MUMFORD, W.W., & RABINOWITZ, J.R. (1980) The increased passive efflux of sodium and rubidium from rabbit erythrocytes by microwave radiation. *Radiat. Res.*, **82**: 244-256.
- OLSEN, R.G. (1982) Far-field dosimetric measurements in a full-sized man model at 2.0 GHz. *Bioelectromagnetics*, **3**: 433-441.
- OLSEN, R.G. & GRINER, T.A. (1982) Electromagnetic dosimetry in a sitting rhesus model at 225 MHz. *Bioelectromagnetics*, **3**: 385-389.
- ONORM (1986) [Microwave and radiofrequency electromagnetic fields; definitions, limits of exposure, measurements.] Vienna, Österreichisches Normungsinstitut (Onorm S1120) (in German).
- OSCAR, K.J. & HAWKINS, T.D. (1977) Microwave alteration of the blood-brain barrier system of rats. *Brain Res.*, **126**: 281-293.
- OSEPCHUK, J.M. (1979) A review of microwave oven safety. *Microwave J.*, **22**: 25-37.
- PARKER, L.N. (1973) Thyroid suppression and adrenomedullary activation by low-intensity microwave radiation. *Am. J. Physiol.*, **224**: 1388-1390.
- PENNES, H. H. (1948) Analysis of tissue and arterial blood temperatures in the resting human forearm. *J. appl. Physiol.*, **1**: 93-122.
- PEREZ, C.A., PAJAK, T.F., EMAMI, B.M., HORNBACH, N.B., TUPCHONG, L., & RUBIN, P. (1991) Randomized phase - III. Study comparing irradiation and hyperthermia with irradiation alone in superficial measurable tumours: final report by the Radiation Therapy Oncology Group. *Am. J. clin. Oncol. Cancer Clin. Trials (USA)* **14**(2): 133-141.
- PETROVICK, Z., LANGHOLZ, B., GIBBS, F.A., SAPOZINK, M.D., KAPP, D.S., STEWART, R.J., EMAMI, B., OLESON, J., SENZER, N., SLATER, J., & ASTRAHAN, M. (1989) Regional hyperthermia for advanced tumours: a clinical study of 353 patients. *Int. J. Radiat. Oncol. Biol. Phys.*, **16**: 601-607.

EHC 137: Electromagnetic fields

- PHILLIPS, R.D., HUNT, E.L., CASTRO, R.D., & KING, N.W. (1975) Thermoregulatory, metabolic and cardiovascular responses of rats to microwaves. *J. appl. Physiol.*, **38**: 630-635.
- POLK, C. & POSTOW, E., ed. (1986) CRC handbook of biological effects of electromagnetic fields. Boca Raton, Florida, CRC Press.
- POLSON, P. & MERRITT, J.H. (1985) Cancer mortality and Air Force bases: A reevaluation. *J. Bioelec.*, **4**: 121-127.
- PRATO, F.S., FRAPPIER, R.H., SHIVERS, R.R., KANKIERS, M., ZABEL, P., DROST, D.J., & LEE, T.Y. (1990) Magnetic resonance imaging increases the brain space of 153 gadolinium diethylene tri-aminopentascetic acid in rats. In: Abstracts, 12th Annual Meeting of the Bioelectromagnetic Society, June 1990, San Antonio, Texas, p. 46.
- PRÄUSNITZ, S. & SUSSKIND, C. (1962) Effects of chronic microwave irradiation on mice. *IRE Trans. Biomed. Electron.*, **9**: 104-108.
- PRESKORN, S.H., EDWARDS, W.D., & JUSTESEN, D.R. (1978) Retarded tumor growth and greater longevity in mice after fetal irradiation by 2450 MHz microwaves. *J. Surg. Oncol.*, **10**: 483-492.
- PRINCE, J. E., MORI, L.H., FRAZER, J.W., & MITCHELL, J.C. (1972) Cytologic aspect of RF radiation in the monkey. *Aerosp. Med.*, **43**: 759-761.
- RAGAN, H.A., PHILLIPS, R.D., BUSCHBOM, R.L., BUSCH, R.H., & MORRIS, J.E. (1983) Haematologic and immunologic effects of pulsed microwaves in mice. *Bioelectromagnetics*, **4**: 383-396.
- RAMA RAO, G., CAIN, C.A., LOCKWOOD, J., & TOMPKINS, W.A.F. (1983) Effects of microwave exposure on the hamster immune system. II. Peritoneal macrophage function. *Bioelectromagnetics*, **4**: 141-155.
- RAMA RAO, G., CAIN, C.A., & TOMPKINS, W.A.F. (1985) Effects of microwave exposure on the hamster immune system. IV. Spleen cell IgM haemolytic plaque formation. *Bioelectromagnetics*, **6**: 41-52.
- REILLY, J.P. (1988) Electrical models for neural excitation studies. Johns Hopkins University, Applied Physics Laboratory, Tech. Digest, **9**: 44-59.
- REPACHOLI, M.H. (1983a) Sources and applications of radiofrequency and microwave energy. In: Grandolfo, M., Michaelson, S.M., & Rindi, R., ed. Biological effects and dosimetry of nonionizing radiation:

radiofrequency and microwave energies. New York, London, Plenum Press, pp. 19-41.

REPACHOLI, M.H. (1983b) Development of standards - Assessment of health hazards and other factors. In: Grandolfo, M., Michaelson, S.M., & Rindi, A., ed. Biological effects and dosimetry of nonionizing radiation: radiofrequency and microwave energies. New York, London, Plenum Press, pp. 611-625.

REPACHOLI, M.H. (1985) Video display terminals - should operators be concerned? Austral. phys. engin. Sci. Med., 8(2): 51-61.

REPACHOLI, M.H., ed. (1988) Non-ionizing radiations: physical characteristics, biological effects and health hazard assessment. London, IRPA Publications, 464 pp.

REPACHOLI, M.H. (1990) Radiofrequency field exposure standards: Current limits and the relevant bioeffects data. In: Gandhi, O.P., ed. Biological effects and medical applications of electromagnetic fields. Englewood Cliffs, New Jersey, Prentice Hall, pp. 9-27.

RHEE, K.W., LEE, C.S., DAVIS, C.C., SAGRIPANTI, J.L., & SWICORD, M.L. (1988) Further studies of the microwave absorption characteristics of different forms of DNA in solution. (Abstract). 10th Annual Meeting of Bioelectromagnetics Society, Stamford, Connecticut, p. 17.

ROBERTI, B., HEEBELS, G.H., HENDRICX, J.C.M., DE GREEF, A.H.A.M., & WOLTHUIS, O.L. (1975) Preliminary investigations of the effects of low-level microwave radiation on spontaneous motor activity in rats. Ann. N.Y. Acad. Sci., 247: 417-424.

ROBERTS, N.J., Jr (1979) Temperature and host defence. Microbiol. Rev., 43: 241-259.

ROBERTS, N.J., Jr (1983) Radiofrequency and microwave effects on immunological and haematopoietic systems. In: Grandolfo, M., Michaelson, S.M., & Rindi, A., ed. Biological effects and dosimetry of nonionizing radiation, radiofrequency and microwave energies. New York, London, Plenum Press, pp. 429-459.

ROBERTS, N.J., Jr, LU, S.T., & MICHAELSON, S.M. (1983) Human leukocyte functions and the US safety standard for exposure to radiofrequency radiation. Science, 220: 318-320.

ROBERTS, N.J., Jr, MICHAELSON, S.M., & LU, S.T. (1984) Exposure of human mononuclear leukocytes to microwave energy pulse modulated at 16 or 60 Hz. IEEE Trans. microwave Theory Tech., MTT-32: 803-807.

ROBERTS, N.J., Jr, MICHAELSON, S.M., & LU, S.T. (1986) The biological effects of radiofrequency radiation: A critical review and recommendations. Int. J. radiat. Biol., 50: 379-420.

ROBINETTE, C.D. & SILVERMAN, C. (1977) Causes of death following occupational exposure to microwave radiation (radar) 1950-1974. In: Hazzard, D.G., ed. Symposium on the Biological Effects and Measurement of Radiofrequency/Microwaves. Washington, DC, Department of Health, Education, and Welfare (HEW Publication No (FDA) 77-8026).

ROBINETTE, C.D., SILVERMAN, C., & JABLON, S. (1980) Effects upon health of occupational exposure to microwave radiation (radar). Am. J. Epidemiol., 112: 39-53.

ROGERS, S.J. (1981) Radiofrequency burn hazards in the MF/HF band. (Aeromedical Review 3-81), pp. 76-89. In: Proceedings of a Workshop on the Protection of Personnel Against Radiofrequency Electromagnetic Radiation, Texas, Brooks Air Force Base, USAF/SAM Aerospace Medical Division.

ROSENTHAL, S.W., BIRENBAUM, L., KAPLAN, I.T., METLAY, W., SNYDER, W.Z., & ZARET, M.M. (1976) Effects of 35 and 107 GHz CW microwaves on the rabbit eye. In: Johnson, C.C. & Shore, M.L., ed. Biological effects of electromagnetic waves. Selected Papers of the USNC/URSI Annual Meeting, Boulder, Colorado, October 1975. Rockville, Maryland, US Department of Health, Education, and Welfare, Vol. 1, pp. 110-128 (HEW Publication (FDA) 77-8010).

ROSS, S.M., LIBURDY, R.P., BUDINGER, T.F., SALFORD, L.S., BRUN, A., PERSSON, B.R.R., ROOS, M.S., de MARINCOR, O.J., & BRENNAN, K.M. (1990) Possibility that the blood-bone barrier (BBB) of the rat to albumin is not significantly altered by nuclear magnetic resonance imaging (NMRI) fields. In: Abstracts, 12th Annual Meeting of the Bioelectromagnetics Society, June, 1990, San Antonio, Texas. p. 46.

→ ROSZKOWSKI, W., WREMBEL, J.K., ROSZKOWSKI, K., JANIAK, M., & SZMIGIELSKI, S. (1980) Does whole-body hyperthermia therapy involve participation of the immune system? Int. J. Cancer, 25: 289-292.

References

ROTKOVSKA, D., VACEK, A., & BARTONICKOVA, A. (1987) Effects of microwaves on the colony-forming ability of haemopoietic stem cells in mice. *Acta oncol.*, 26: 233-236.

ROZZELL, T.C. (1985) West Germany EMF exposure standard (BEMS Newsletter, 55).

RUGGERA, P.S. (1980) Measurements of emission levels during microwave and short wave diathermy treatments. Rockville, Maryland, US Department of Health and Human Services, FDA (Publication No. FDA 80-8119).

SAA (1988) Radio-frequency radiation - principles and methods of measurement. Sydney, Standards Association of Australia.

SAGER, D.P. (1987) Current facts on pacemaker electromagnetic interference and their application to clinical care. *Heart Lung*, 16: 211-221.

SANDSTROM, M., HANSSON-MILD, K., & LOVTRUP, S. (1987) Effects of weak pulsed magnetic fields on chick embryogenesis. In: Knave, B. & Wideback, P.G., ed. *Work with display units 86*. Amsterdam, Elsevier, p. 135.

SANTINI, R., HOSNI, M., DESCHAUX, P., & PACKECO, H. (1988) B16 melanoma development in black mice exposed to low-level microwave radiation. *Bioelectromagnetics*, 9(1): 105-107.

SANZA, J.N. & DE LORGE, J. (1977) Fixed interval behaviour of rats exposed to microwaves at low power densities. *Radio Sci.*, 12: 273-277.

SAUNDERS, R.D., & KOWALCZUK, C.I. (1981) Effects of 2.45 GHz microwave radiation and heat on mouse spermatogenic epithelium. *Int. J. radiat. Biol.*, 40: 623-632.

SAUNDERS, R.D., DARBY, S.C., & KOWALCZUK, C.I. (1983) Dominant lethal studies in male mice after exposure to 2.45 GHz microwave radiation. *Mutat. Res.*, 117: 345-356.

SAUNDERS, R.D., KOWALCZUK, C.I., BEECHEY, C.V., & DUNFORD, R. (1988) Studies of the induction of dominant lethals and translocations in male mice after chronic exposure to microwave radiation. *Int. J. radiat. Biol.*, 53: 983-992.

SAUNDERS, R.D., KOWALCZUK, C.I., & SIENKIEWICZ, Z.J. (1991) The biological effects of non-ionizing electromagnetic fields and radiation:

EHC 137: Electromagnetic fields

III. Radiofrequency and microwave radiation. Oxfordshire, England, National Radiological Protection Board (NRPB R 240).

SAVIN, B.M. (1986) Safety regulations for non-ionizing radiation. In: *Hygienic standardization of NIR*. Moscow, Medicina, pp. 115-146.

SAVIN, B.M., NIKONOVA, K.W., LOBANOVA, E.A., SADCZIKOVA, M.N., & LOBED, E.K. (1983) [Novelties in safety standards of EM radiation of the microwave range.] *Gig. Truda*, 3: 1 (in Russian).

SCHAEFER, D.J., BARBER, B.J., GORDON, C.J., ZIELONKA, J. & HECKER, J. (1985) Thermal effects of magnetic resonance imaging (MRI). In: *Abstracts, Meeting of the Society of Magnetic Resonance in Medicine*, Vol. 2, pp. 925-926, Berkeley, California, Society of Magnetic Resonance in Medicine.

SCHLAGEL, C.J. & AHMED, A. (1982) Evidence for genetic control of microwave-induced augmentation of complement receptor-bearing B lymphocytes. *J. Immunol.*, 129(4): 1530-1533.

SCHLAGEL, C.J., SULEK, K., HO, H.S., LEACH, W.M., AHMED, A., & WOODY, J.N. (1980) Biological effects of microwave exposure. II Studies on the mechanisms controlling susceptibility to microwave-induced increases in complement receptor-positive spleen cells. *Bioelectromagnetics*, 1: 405-414.

SCHNORR, T.M., GRAJEWSKI, B.A., HORNING, R.W., THUN, M.J., EGELAND, G.M., MURRAY, W.E., CONOVER, D.L., & HALPERIN, W.E. (1991) Video display terminals and the risk of spontaneous abortion. *New England J. Med.*, 324: 727-733.

SCHOLL, D.J. & ALLEN, S.J., (1979) Skilled visual-motor performance by monkeys in a 1.2-GHz microwave field. *Radio Sci.*, 12: 247-252.

SCHROT, J., THOMAS, J.R., & BANVARD, R.A. (1980) Modification of the repeated acquisition of response sequences in rats by low-level microwave exposure. *Bioelectromagnetics*, 1: 89-99.

SCHWAN, H.P. (1984) Frequency selective propagation of extracellular electrical stimuli to intracellular compartments. In: Adey, W.R. & Lawrence, A.F., ed. *Nonlinear electrodynamics in biological systems*. New York, London, Plenum Press, pp. 327-338.

SCHWAN, H.P. (1985) Biophysical principles of interactions and forces. In: Grandolfo, M., Michaelson, S.M., & Rindi, A., ed. *Biological effects*

References

- STUCHLY, M.A., SPIEGEL, R.J., STUCHLY, S.S., & KRASZEWSKI, A. (1986) Exposure of man in the near-field of a resonant dipole: comparison between theory and measurements. *IEEE Trans. microwave Theory Tech.*, **MTT-34**: 26-30.
- STUCHLY, M.A., KRASZEWSKI, A., STUCHLY, S.S., HARTSGROVE, G.W., & SPIEGEL, R.J. (1987) Energy deposition in a heterogeneous model of man: near-field exposures. *IEEE Trans. Biomed. Eng.*, **BME-34**: 944-950.
- STUCHLY, M.A., RUDDICK, J., VILLENEUVE, D., ROBINSON, K., REED, B., LECUYER, D.W., TAN, K., & WONG, J. (1988) Teratological assessment of exposure to time-varying magnetic field. *Teratology*, **38**: 461.
- STUCHLY, S.S., KRASZEWSKI, A., STUCHLY, M.A., HARTSGROVE, G., & ADAMSKI, D. (1985) Energy deposition in a model of man in the near-field. *Bioelectromagnetics*, **6**: 115-129.
- STUCHLY, S.S., STUCHLY, M.A., KRASZEWSKI, A., & HARTSGROVE, G. (1986) Energy deposition in a model of man; frequency effects. *IEEE Trans. biomed. Eng.*, **BME-33**: 702-711.
- STUCHLY, S.S., KRASZEWSKI, A., STUCHLY, M.A., HARTSGROVE, G., & SPIEGEL, R.J. (1987) RF energy deposition in a heterogeneous model of man: far-field exposures. *IEEE Trans. biomed. Eng.*, **BME-34**: 951-957.
- SUESS, M.J. & BENWELL-MORISON, D.A., ed. (1989) Non-ionizing radiation protection, 2nd ed. Copenhagen, World Health Organization Regional Office for Europe, 346 pp. (European Series No. 25).
- SULTAN, M.F., CAIN, C.A., & TOMPKINS, W.A.F. (1983a) Effects of microwaves and hyperthermia on capping of antigen-antibody complexes on the surface of normal mouse B lymphocytes. *Bioelectromagnetics*, **4**: 115-122.
- SULTAN, M.F., CAIN, C.A., & TOMPKINS, W.A.F. (1983b) Immunological effects of amplitude-modulated radiofrequency radiation: B lymphocyte capping. *Bioelectromagnetics*, **4**: 157-166.
- SZMIGIELSKI, S. & OBARA, T. (1989) The rationale for the Eastern European radiofrequency and microwave protection guides. In: Franceschetti, G., Gandhi O.P., & Grandolfo, M., ed. *Electromagnetic biointeraction - Mechanisms, safety standards, protection guides*. New York, London, Plenum Press, pp. 135-151.
- SZMIGIELSKI, S., SZUDZINSKI, A., PIETRASZEK, A., BIELEC, M., & WREMBEL, J.K. (1982) Accelerated development of spontaneous and benzo(a)pyrene-induced skin cancer in mice exposed to 2450 MHz microwave radiation. *Bioelectromagnetics*, **3**: 179-191.
- SZMIGIELSKI, S., BIELEC, M., LIPSKI, S., & SOKOLSKA, G. (1988) Immunologic and cancer-related aspects of exposure to low-level microwave and radiofrequency fields. In: Marino, A.A., ed. *Modern bioelectricity*. New York, Marcel Dekker, Inc., pp. 861-925.
- SZUDZINSKI, A., PIETRASZEK, A., JANIAK, M., WREMBEL, J., KALCZEK, M., & SZMIGIELSKI, S. (1982) Acceleration of the development of benzo(a)pyrene-induced skin cancer in mice by microwave radiation. *Arch. dermatol. Res.*, **274**: 303-312.
- TAKASHIMA, S., ONARAL, B., & SCHWAN, H.P. (1979) Effects of modulated RF energy on the EEG of mammalian brains. *Radiat. environ. Biophys.*, **16**: 15-27.
- TAKASHIMA, S., GABRIEL, C., SHEPPARD, R.J., & GRANT, E.H. (1984) Dielectric behaviour of DNA solutions at radio and microwave frequencies (at 20 °C). *Biophys. J.*, **46**: 29-34.
- TELL, R.A. (1983) Instrumentation and measurement of electromagnetic fields: Advanced Study Institute, series A. *Life Sci.*, **49**: 95-162.
- TELL, R.A. (1990) RF hot spot fields: The problem of determining compliance with the ANSI radiofrequency protection guide. *NAB Engineering Conference Proceedings*, pp. 419-431.
- TELL, R. A. & MANTIPLY, E. D. (1980) Population exposure to VHF and UHF broadcast radiation in the United States. *Proc. IEEE*, **68**: 6-12.
- TELL, R.A., MANTIPLY, E.D., DURNEY, C.H., & MASSOUDI, H. (1982) Electric and magnetic field intensities and associated induced body currents in man in close proximity to a 50 kW AM standard broadcast station. Las Vegas, Nevada, US Environmental Protection Agency, Electromagnetic Radiation Analysis Branch, and Salt Lake City, Utah, Departments of Electrical Engineering and Bioengineering, University of Utah.